



POSTER PRESENTATION

Open Access

Quantitative texture features as objective metrics of enhancement heterogeneity in hypertrophic cardiomyopathy

Rebecca E Thornhill^{1,2*}, Myra Cocker³, Girish Dwivedi³, Carole Dennie^{1,2}, Lyanne Fuller³, Alexander Dick³, Terrence Ruddy³, Elena Pena^{1,2}

From 17th Annual SCMR Scientific Sessions
New Orleans, LA, USA. 16-19 January 2014

Background

Hypertrophic cardiomyopathy (HCM) results in myocardial disarray, hypertrophy and fibrosis. Late gadolinium enhanced MRI (LGE) can assess the presence and extent of fibrosis, which is associated with the development of arrhythmias and sudden cardiac death. However, enhancement may not always be present or only sparsely distributed. Thus, one of the challenges is how best to describe heterogeneous LGE patterns in an objective fashion that informs clinical decision making. Quantitative texture features may provide clinicians with an objective means of describing the heterogeneity of LGE patterns in HCM. We hypothesized that hypertrophied segments would exhibit greater grey-level heterogeneity than both (a) non-hypertrophied segments in HCM patients, and (b) healthy volunteers.

Methods

We prospectively recruited 12 HCM patients and 4 healthy volunteers. Functional (bSSFP cine) and LGE (phase-sensitive inversion recovery spoiled GRE, 10-15 min post injection of 0.2 mmol/kg Gd-DTPA) images were acquired in short-axis orientation (SAO), as well as in one 4-chamber slice. We measured the maximum thickness on the end-diastolic cine frame in the 17 segments (AHA model). Segments measuring > 15 mm on SSFP images were considered hypertrophic (H+). Segments were categorized as fibrotic (F+) if > 20% of pixels were enhanced (> 5 SD nulled myocardium). The extent of myocardial fibrosis on LGE imaging and textural features (run-length non-uniformity, RLNU, and

grey-level non-uniformity, GLNU [Galloway 1975]) were assessed for each segment. Differences in RLNU and GLNU among segment groups (H+/F+, H+/F-, H-/F+, H-/F-, and healthy) were assessed by Kruskal-Wallis tests.

Results

Of 192 segments we found; 7 H+/F+, 9 H+/F-, 29 H-/F+, and 147 H-/F-. Median +/-interquartile ranges for RLNU and GLNU for each HCM group, as well as for the 64 segments obtained from healthy volunteers are depicted in Figure 1 ($P < 0.0001$, for RLNU and GLNU). Post-hoc analysis revealed that RLNU and GLNU were significantly greater in H+ than H- segments ($P = 0.006$ and $P = 0.0002$, respectively). Both RLNU and GLNU in H-/F- HCM segments were greater than in healthy volunteers ($P = 0.009$ and $P < 0.0001$, respectively).


Conclusions

Quantitative textural features related to LGE heterogeneity appear elevated in patients with HCM, even in non-hypertrophic segments. In addition, significant statistical differences were found in the textural features between non-hypertrophic, non-fibrotic segments of HCM patients and healthy volunteers. Thus, RLNU and GLNU show potential for markers of incipient cardiomyopathic changes among HCM patients and may provide helpful tools for differentiating diverse phenotypic expressions of the disease from healthy patients, pending further validation.

Funding

Nothing to disclose.

¹Medical Imaging, The Ottawa Hospital, Ottawa, Ontario, Canada
Full list of author information is available at the end of the article



Authors' details

¹Medical Imaging, The Ottawa Hospital, Ottawa, Ontario, Canada. ²Radiology, University of Ottawa, Ottawa, Ontario, Canada. ³Cardiology, University of Ottawa Heart Institute, Ottawa, Ontario, Canada.

Published: 16 January 2014

doi:10.1186/1532-429X-16-S1-P351

Cite this article as: Thornhill *et al.*: Quantitative texture features as objective metrics of enhancement heterogeneity in hypertrophic cardiomyopathy. *Journal of Cardiovascular Magnetic Resonance* 2014 **16** (Suppl 1):P351.

Submit your next manuscript to BioMed Central and take full advantage of:

- Convenient online submission
- Thorough peer review
- No space constraints or color figure charges
- Immediate publication on acceptance
- Inclusion in PubMed, CAS, Scopus and Google Scholar
- Research which is freely available for redistribution

Submit your manuscript at
www.biomedcentral.com/submit

